arXiv:1305.0820v1 [gr-qc] 3 May 2013

Dark matter density profile and galactic metric in Eddington-inspired Born-Infeld gravity

Tiberiu Harko[1],[*] Francisco S. N. Lobo[2],[†] M. K. Mak[3],[‡] and Sergey V. Sushkov[4§]
[1]*Department of Mathematics, University College London,*
Gower Street, London WC1E 6BT, United Kingdom
[2]*Centro de Astronomia e Astrofísica da Universidade de Lisboa,*
Campo Grande, Ed. C8 1749-016 Lisboa, Portugal
[3]*Department of Computing and Information Management,*
Hong Kong Institute of Vocational Education, Chai Wan, Hong Kong, and
[4]*Institute of Physics, Kazan Federal University, Kremlevskaya Street 18, Kazan 420008, Russia*
(Dated: May 7, 2013)

We consider the density profile of pressureless dark matter in Eddington-inspired Born-Infeld (EiBI) gravity. The gravitational field equations are investigated for a spherically symmetric dark matter galactic halo, by adopting a phenomenological tangential velocity profile for test particles moving in stable circular orbits around the galactic center. The density profile and the general form of the metric tensor is obtained in an analytical form for both physical-g and auxiliary-q metrics, to a first order approximation. The dark matter density distribution is described by the Lane-Emden equation with polytropic index $n = 1$, and is non-singular at the galactic center. The gravitational properties of the dark matter halos are also briefly discussed in the Newtonian approximation.

Introduction. Despite numerous observational and experimental attempts, up to now the elusive dark matter particles have evaded detection. Great hopes in gaining more insights into the nature of dark matter are related to the Alpha Magnetic Spectrometer (AMS-02) collaboration results, which have just been released [1]. The AMS experiment has determined the positron fraction in cosmic rays with extremely high precision. The positron fraction rises continuously from ~ 5 GeV up to ~ 350 GeV, while the slope becomes flat above ~ 100 GeV [1]. In order to explain the AMS results, dark matter is still an attractive candidate. In fact, dark matter annihilation into $\tau^+\tau^-$ final states, which results in a soft positron spectrum can account for the AMS-02 data quite well. Other channels for dark matter annihilation/decay into multiple μ or τ leptons can also reproduce the AMS-02 data [2]. However, presently, the only *convincing* evidence, based on the flat rotation galactic curves and the virial mass discrepancy in clusters of galaxies for the existence of dark matter, is *gravitational* [3].

In standard general relativity the coupling between matter and gravity is given by a proportionality relation between the stress-energy tensor and the geometry. Although both the stress-energy tensor and the Einstein tensor are divergenceless, there is no obvious reason why the matter-gravity coupling should be linear [4]. On the other hand, modified theories of gravity usually affect the vacuum dynamics, yet keep the matter-gravity coupling linear. Recently, to provide matter-gravity coupling modifications, based on the Eddington gravity action [5], and Born-Infeld nonlinear electrodynamics [6], the Eddington-inspired Born-Infeld (EiBI) theory has been proposed [7]. The EiBI theory coupled to a perfect fluid reduces to standard general relativity coupled to a nonlinearly modified perfect fluid, leads to an ambiguity between modified coupling and a modified equation of state [8, 9]. The theory has interesting cosmological consequences, leading to the possibility of avoiding cosmological singularities [10]. However, the EiBI theory is reminiscent of the Palatini gravity formulation, and it shares the same pathologies, such as curvature singularities at the surface of polytropic stars [11].

It is the purpose of this Letter to investigate the properties of dark matter halos in the EiBI theory. By assuming a spherically symmetric pressureless dark matter halo, the gravitational field equations of the EiBI model are solved to a first order approximation of the coupling constant κ. The dark matter density profile and the galactic metric in the dark matter halo is explored, and the respective gravitational properties are also briefly discussed in the Newtonian approximation.

Eddington-inspired Born-Infeld gravity. The starting point of the EiBI theory is the gravitational action S, given by [7],

$$S = \frac{1}{16\pi}\frac{2}{\kappa}\int d^4x\left(\sqrt{-\left|g_{\mu\nu}+8\pi\kappa R_{\mu\nu}\right|}-\lambda\sqrt{-g}\right) + S_M\left[g,\Psi_M\right], \tag{1}$$

where $\lambda \neq 0$ is a dimensionless parameter, and κ is a parameter with inverse dimension to that of the cosmological constant Λ. $R_{\mu\nu}$ is the symmetric part of the Ricci tensor, and is constructed solely from the connection $\Gamma^\alpha_{\beta\gamma}$. The matter action S_M depends only on the metric $g_{\mu\nu}$ and the matter fields Ψ_M. The determinant of the tensor $g_{\mu\nu} + 8\pi\kappa R_{\mu\nu}$ is denoted by $|g_{\mu\nu} + 8\pi\kappa R_{\mu\nu}|$. In the limit $\kappa \to 0$, the action Eq. (1) recovers the Einstein-Hilbert action with $\lambda = 8\pi\Lambda\kappa + 1$. However, in the present paper, we consider only asymptotic flat solutions, and hence take $\lambda = 1$. Therefore the cosmological constant vanishes, and the remaining parameter κ plays the determining role for the description of the physical behavior of various cosmological and stellar scenarios. Several constraints on the value and the sign of the parameter κ have been obtained from solar observations, big bang nucleosynthesis, and the existence of neutron stars in [7, 12]. The structure of compact stars in EiBI theory has been investigated by several authors [12, 13]. In particular, for cases with positive κ, effective gravitational repulsion prevails, leading to the existence of pressureless stars (stars made of non-interacting particles which provide interesting models for self-gravitating dark matter) and increases in the mass limits of compact stars [12, 13].

---

[*] Electronic address: t.harko@ucl.ac.uk
[†] Electronic address: flobo@cii.fc.ul.pt
[‡] Electronic address: mkmak@vtc.edu.hk
[§] Electronic address: sergey˙sushkov@mail.ru

In the EiBI theory the metric $g_{\mu\nu}$ and the connection $\Gamma^{\alpha}_{\beta\gamma}$ are treated as independent fields. By varying the action (1) with respect to the connection $\Gamma^{\alpha}_{\beta\gamma}$, and with respect to the real metric $g_{\mu\nu}$, yield the following field equations $q_{\mu\nu} = g_{\mu\nu} + 8\pi\kappa R_{\mu\nu}$ and $q^{\mu\nu} = \tau\left(g^{\mu\nu} + 8\pi\kappa T^{\mu\nu}\right)$, respectively. The auxiliary metric $q_{\mu\nu}$ is related to the connection by $\Gamma^{\alpha}_{\beta\gamma} = \frac{1}{2}q^{\alpha\sigma}\left(\partial_\gamma q_{\sigma\beta} + \partial_\beta q_{\sigma\gamma} - \partial_\sigma q_{\beta\gamma}\right)$, and τ is defined as $\tau = \sqrt{g/q}$. If the stress-energy tensor $T^{\mu\nu}$ vanishes, then the real metric $g_{\mu\nu}$ is equal to the apparent metric $q_{\mu\nu}$. Hence, in vacuum the EiBI theory is equivalent to standard general relativity. The stress-energy tensor $T^{\mu\nu}$ satisfies the conservation equations given by $\nabla_\mu T^{\mu\nu} = 0$, where the covariant derivative ∇_μ refers to the metric $g_{\mu\nu}$.

Static spherically symmetric dark matter halos in EiBI gravity. We consider that the dark matter halo is static and spherically symmetric. Therefore the line elements for the physical metric $g_{\mu\nu}$ and for the auxiliary metric $q_{\mu\nu}$ take the forms

$$
\begin{aligned}
g_{\mu\nu}dx^\mu dx^\nu &= -e^{\nu(r)}dt^2 + e^{\lambda(r)}dr^2 + f(r)\,d\Omega^2, && (2)\\
q_{\mu\nu}dx^\mu dx^\nu &= -e^{\beta(r)}dt^2 + e^{\alpha(r)}dr^2 + r^2 d\Omega^2, && (3)
\end{aligned}
$$

respectively, where $\nu(r)$, $\lambda(r)$, $\beta(r)$, $\alpha(r)$ and $f(r)$ are arbitrary metric functions of the radial coordinate r, and $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$. The galactic halo is made up of a perfect fluid described by the standard stress-energy tensor $T^{\mu\nu} = (\rho + p)\,u^\mu u^\nu + pg^{\mu\nu}$, where ρ, p and u^μ are the energy density, the isotropic pressure and the four velocity of the fluid, respectively, with the latter satisfying $u^\mu u^\nu g_{\mu\nu} = -1$. The system of gravitational field equations describing the dark matter halo structure yields [13]

$$\frac{d}{dr}\left(re^{-\alpha}\right) = 1 - \frac{1}{2\kappa}\left(2 + \frac{a}{b^3} - \frac{3}{ab}\right)r^2, \qquad (4)$$

$$e^{-\alpha}\left(1 + r\frac{d\beta}{dr}\right) = 1 + \frac{1}{2\kappa}\left(\frac{1}{ab} + \frac{a}{b^3} - 2\right)r^2, \qquad (5)$$

$$e^{\beta} = \frac{e^{\nu}b^3}{a}, \qquad e^{\alpha} = e^{\lambda}ab, \qquad f = \frac{r^2}{ab}, \qquad (6)$$

where we have defined the arbitrary functions $a(r)$ and $b(r)$ as $a = \sqrt{1 + 8\pi\kappa\rho}$, and $b = \sqrt{1 - 8\pi\kappa p}$, respectively. In the g–metric the conservation of the stress-energy tensor yields the result

$$\frac{d\nu}{dr} = -\frac{2}{p + \rho}\frac{dp}{dr} = \frac{4b}{a^2 - b^2}\frac{db}{dr}. \qquad (7)$$

The existence of dark matter at the galactic scale is inferred from the study of the rotational velocities of massive test particles (hydrogen clouds) around the galactic center. The Lagrangian $\mathcal{L}$ for a massive test particle reads $\mathcal{L} = \frac{1}{2}(-e^{\nu}\dot{t}^2 + e^{\lambda}\dot{r}^2 + r^2\dot{\Omega}^2)$, where the overdot denotes differentiation with respect to the affine parameter s. By defining the tangential velocity $v_{\rm tg}$ of a test particle, as measured in terms of the proper time [14], that is, by an observer located at a given point, as $v_{tg}^2 = e^{-\nu}r^2\left(d\Omega/dt\right)^2$, we obtain the expression of v_{tg}^2 for a test particle in a stable circular orbit as $v_{tg}^2 = r\nu'/2$ [15].

For v_{tg}^2 we assume the simple empirical dark halo rotational velocity law [16]

$$v_{tg}^2 = v_\infty^2\frac{\left(r/r_{opt}\right)^2}{\left(r/r_{opt}\right)^2 + r_0^2}, \qquad (8)$$

where r_{opt} is the optical radius containing 83% of the galactic luminosity. The parameter r_0, defined as the ratio of the halo core radius and r_{opt}, and the terminal velocity v_∞ are functions of the galactic luminosity L. For spiral galaxies $r_0 = 1.5\left(L/L_*\right)^{1/5}$ and $v_\infty^2 = v_{opt}^2\left(1 - \beta_*\right)\left(1 + r_0^2\right)$, where $v_{opt} = v_{tg}\left(r_{opt}\right)$, and $\beta_* = 0.72 + 0.44\log_{10}\left(L/L_*\right)$, with $L_* = 10^{10.4}L_\odot$. For $r \to 0$, $v_{tg} \to 0$, while for $r/r_{opt} \gg r_0$, $v_{tg} \to v_\infty$.

Therefore, the most general static and spherically symmetric metric of the dark matter halo can be written as

$$ds^2 = -e^{\nu_0}\left[\left(\frac{r}{r_{opt}}\right)^2 + r_0^2\right]^{v_\infty^2}dt^2 + e^{\lambda(r)}dr^2 + r^2 d\Omega^2, \qquad (9)$$

where e^{ν_0} is an arbitrary constant of integration.

The dark matter density profile and the galactic metric in EiBI gravity. In order to study the extra-galactic motion of massive test particles in EiBI gravity, we assume that the motion takes place in a pressureless cosmic medium, $8\pi\kappa p \ll 1$, that can be described by an effective matter density ρ. Since $p \ll 1/8\pi\kappa$, we obtain $b \approx 1$. For a galactic metric of the form given by Eq. (9) we obtain

$$
\begin{aligned}
e^{\beta} &= \frac{e^{\nu_0}\left[\left(r/r_{opt}\right)^2 + r_0^2\right]^{v_\infty^2}}{\sqrt{1 + 8\pi\kappa\rho}}, && (10)\\
\beta' &= 2v_\infty^2\frac{r/r_{opt}^2}{\left(r/r_{opt}\right)^2 + r_0^2} - 4\pi\kappa\frac{\rho'}{1 + 8\pi\kappa\rho}, && (11)
\end{aligned}
$$

respectively. In the following, we consider that the density of the dark matter is small, so that $8\pi\kappa\rho \ll 1$. Then, by using a series expansion of a and β' to first order of κ, we obtain $1/\sqrt{1 + 8\pi\kappa\rho} + \sqrt{1 + 8\pi\kappa\rho} - 2 \approx 0$, $2 + \sqrt{1 + 8\pi\kappa\rho} - 3/\sqrt{1 + 8\pi\kappa\rho} \approx 16\pi\kappa\rho$, and $\beta' =$

$2v_\infty^2(r/r_{opt}^2)/[(r/r_{opt})^2+r_0^2]-4\pi\kappa\rho'$. Therefore the field equations Eqs. (4)-(5) become respectively

$$\frac{d}{dr}\left(re^{-\alpha}\right)=1-8\pi\rho r^2, \tag{12}$$

$$re^{-\alpha}=\frac{1}{1/r+2v_\infty^2\left(r/r_{opt}^2\right)/\left[\left(r/r_{opt}\right)^2+r_0^2\right]-4\pi\kappa\rho'} \tag{13}$$

For $r\rightarrow 0$, $1/r\gg 2v_\infty^2\left(r/r_{opt}^2\right)/[(r/r_{opt})^2+r_0^2]$, and therefore in this limit the term containing v_∞^2 can be neglected. For $(r/r_{opt})^2\gg r_0^2$, we obtain $1/r+2v_\infty^2\left(r/r_{opt}^2\right)/[(r/r_{opt})^2+r_0^2]\approx\left(1+2v_\infty^2\right)/r\approx 1/r$, since $v_\infty^2\approx 10^{-6}-10^{-8}$ [16]. Therefore, to a very good approximation, in the range $0\leq r\leq\infty$, Eq. (13) can be written as $re^{-\alpha}=1/(1/r-4\pi\kappa\rho')$. By taking the derivative of the latter with respect to r, we obtain the following differential equation describing the dark matter profile in EiBI gravity,

$$\frac{1}{r^2}\frac{d}{dr}\left(r^2\frac{d\rho}{dr}\right)+\frac{2}{\kappa}\rho=16\pi\rho\rho' r. \tag{14}$$

In order to obtain an approximate solution of Eq. (14), in the first order of approximation, and in the linear regime in ρ, we neglect the term $16\pi\rho\rho' r$ on the right hand side of the equation. Therefore we obtain

$$\frac{1}{r^2}\frac{d}{dr}\left(r^2\frac{d\rho^{(0)}}{dr}\right)+\frac{2}{\kappa}\rho^{(0)}=0. \tag{15}$$

Eq. (15) is the Lane-Emden equation for the polytropic index $n=1$ [17], and its non-singular solution at the center is given by

$$\rho^{(0)}(r)=K\frac{\sin\left(\sqrt{2/\kappa}r\right)}{\sqrt{2/\kappa}r}, \tag{16}$$

where K is an arbitrary constant of integration. Since at the center of the galactic halo the density of the dark matter is $\rho^{(0)}(0)=\rho_0$, it follows that $K=\rho_0$. In the EiBI theory the dark halo has a sharp boundary R_{DM}, corresponding to $\rho^{(0)}(R_{DM})=0$, which gives $R_{DM}=\sqrt{\kappa/2}\pi$. Thus, the mass profile of the dark matter $M(r)=4\pi\int_0^r\rho^{(0)}(r)r^2dr$ is given by

$$M(\bar{r})=\frac{4R_{DM}^3}{\pi^2}\rho_0\left[\sin(\bar{r})-\bar{r}\cos(\bar{r})\right], \tag{17}$$

where $\bar{r}=\pi r/R_{DM}$ is defined for convenience.

The q-metric coefficient $e^{-\alpha}=1-2M(r)/r$ yields

$$\begin{aligned} e^{-\alpha} &= 1-\frac{8R_{DM}^2}{\pi\bar{r}}\rho_0\left[\sin(\bar{r})-\bar{r}\cos(\bar{r})\right] \\ &= 1-\frac{\bar{\rho}_0}{\bar{r}}\left[\sin(\bar{r})-\bar{r}\cos(\bar{r})\right], \end{aligned} \tag{18}$$

where $\bar{\rho}_0=8\rho_0R_{DM}^2/\pi$. For the metric coefficient $e^{-\lambda}=e^{-\alpha}a$, we obtain

$$e^{-\lambda}\approx e^{-\alpha}\left(1+4\pi\kappa\rho\right)=e^{-\alpha}\left[1+\frac{\bar{\rho}_0}{\bar{r}}\sin(\bar{r})\right], \tag{19}$$

while the function $f(\bar{r})$ is given by

$$f(\bar{r})=\frac{\left(R_{DM}^2/\pi^2\right)\bar{r}^2}{\sqrt{1+8\pi\kappa\rho}}\approx\frac{R_{DM}^2}{\pi^2}\bar{r}^2\left[1-\frac{\bar{\rho}_0}{\bar{r}}\sin(\bar{r})\right]. \tag{20}$$

Thus, the metric of the dark matter halo takes the form

$$\begin{aligned} ds^2 &= -e^{\nu_0}\left[\left(\frac{R_{DM}}{\pi r_{opt}}\right)^2\bar{r}^2+r_0^2\right]^{v_\infty^2}dt^2 \\ &+ \frac{R_{DM}^2}{\pi^2}\frac{d\bar{r}^2}{\left[1-\frac{\bar{\rho}_0}{\bar{r}}\sin(\bar{r})+\bar{\rho}_0\cos(\bar{r})\right]\left[1-\frac{\bar{\rho}_0}{\bar{r}}\sin(\bar{r})\right]} \\ &+ \frac{R_{DM}^2}{\pi^2}\bar{r}^2\left[1-\frac{\bar{\rho}_0}{\bar{r}}\sin(\bar{r})\right]d\Omega^2. \end{aligned} \tag{21}$$

The total mass of the dark matter halo is given by $M_{DM}=M\left(R_{DM}\right)=\sqrt{2}\pi^2\kappa^{3/2}\rho_0=(4/\pi)\rho_0R_{DM}^3$. The mean density $<\rho>$ of the dark matter halo is obtained as $<\rho>=3M_{DM}/4\pi R_{DM}^3=3\rho_0/\pi^2$.

In the Newtonian approximation the gravitational potential $V_{grav}(r)$ of the dark matter distribution in the EiBI theory is determined for $\bar{r}/\pi\leq 1$ by

$$V_{grav}=\frac{1}{8\pi}\int_r^{R_{DM}}\frac{M\left(r'\right)dr'}{r'^2}=\frac{\bar{\rho}_0}{16\pi\bar{r}}\sin(\bar{r}). \tag{22}$$

At small radii and for $\bar{r}/\pi\leq 1$ the potential behaves as

$$V_{grav}(\bar{r})\approx\frac{\bar{\rho}_0}{16\pi}\left(1-\frac{1}{3!}\bar{r}^2+\frac{1}{5!}\bar{r}^4-\frac{1}{7!}\bar{r}^6\right)+O(\bar{r})^8. \tag{23}$$

The gravitational potential energy $U(r)$ per unit mass and inside radius r of the dark matter halo yields

$$\begin{aligned} U &= -\frac{1}{2}\int_0^r\frac{\rho^{(0)}(r)M(r)}{r}r^2dr=-\frac{R_{DM}^5\rho_0^2}{4\pi^3}\times \\ &\left\{2\bar{r}\left[2+\cos(2\bar{r})\right]-3\sin(2\bar{r})\right\}. \end{aligned} \tag{24}$$

Thus, the total potential energy of the dark matter halo is given by $U\left(R_{DM}\right)=-\left(3\rho_0^2/2\pi^2\right)R_{DM}^5$.

Conclusions. In the present Letter, we have analyzed the dark matter density profiles, and their geometry, in the EiBI gravitational theory. In the approximations of the dark matter pressure satisfying the condition $8\pi\kappa p\ll 1$, of the constant velocity of test particles in circular stable orbits around the galactic center, and to first order of the coupling constant κ, respectively, the dark matter density profile satisfies the Lane-Emden equation with a polytropic index $n=1$. Therefore the dark matter profile can be obtained in an exact analytic form, as well as the full galactic dark matter metric, in both g and q geometries. Similar results for pressureless stellar profiles have been obtained in [9]. In order to obtain a consistent solution of the field equations, the energy conservation equation Eq. (7) must also be satisfied. By assuming that $p\ll\rho$, it follows that the pressure distribution of the dark matter satisfies the equation

$$\frac{dp}{d\bar{r}}=-\frac{\rho_0\left(v_\infty R_{DM}/\pi r_{opt}\right)^2}{\left(R_{DM}/\pi r_{opt}\right)^2\bar{r}^2+r_0^2}\sin(\bar{r}). \tag{25}$$

In the limit of small r the dark matter pressure distribution is given by $p(\bar{r}) = p_c + \frac{v_\infty^2 \rho_0 R_{DM}^2}{\pi^2 r_{opt}^2}\left[\cos\left(\bar{r}\right) - 1\right]$, where $p_c = p(0)$ is the central pressure, while for $r/r_{opt} \gg r_0$ we obtain $p(\bar{r}) \approx \rho_0 v_{tg}^2 \left[\frac{\sin(\bar{r})}{\bar{r}} - \mathrm{Ci}\left(\bar{r}\right)\right]$, where $\mathrm{Ci}(z) = -\int_z^\infty \left[\cos(t)/t\right] dt$ is the cosine integral function. In the non-relativistic regime, the density ρ of the dark matter is given by $\rho = mn$, where n is the particle number density, and m is the dark particle mass. The dark matter pressure p can be obtained as $p = \left\langle \vec{v}^2 \right\rangle \rho/3$, where $\left\langle \vec{v}^2 \right\rangle$ is the average squared velocity of the particle, and $\left\langle \vec{v}^2 \right\rangle/3$ is the one-dimensional velocity dispersion. In the non-relativistic approximation, the velocity dispersion is a constant only for the case of the non-degenerate ideal Maxwell-Boltzmann gas, whose Newtonian analogue is the isothermal sphere. Therefore, in the present model the average velocity dispersion of the dark matter particles in the constant velocity region is

$$\left\langle \vec{v}^2 \right\rangle = 3v_{tg}^2 \left[1 - \frac{\rho_0}{\rho}\mathrm{Ci}\left(\bar{r}\right)\right] = 3v_{tg}^2 \left[1 - \frac{\bar{r}}{\sin\left(\bar{r}\right)}\mathrm{Ci}\left(\bar{r}\right)\right].$$

Note that in the dark matter models, the $n = 1$ polytrope is used to model dark matter in the form of a Bose-Einstein condensate, namely, an assembly of light individual bosons that acquire a repulsive interaction by occupying the same ground energy state [18]. The experimental tests of the Bose-Einstein condensation in terrestrial laboratories for a large class of particles (atoms) and physical systems has given a sound theoretical and experimental support to this hypothesis. However, while in the Bose-Einstein condensate dark matter model quantum effects play the dominant role in determining the density profile, in the EiBI model matter-gravity coupling, and modified geometry, lead to the similar distribution for pressureless dark matter. The non-singular $n = 1$ polytropic density profiles show the presence of an extended core, whose presence is due to the strong interaction between dark matter particles. All the relevant astrophysical quantities can be predicted from the model, and can be directly compared with the corresponding observational parameters (the dark halo mass, the radius of the galaxy, etc), with all the properties of the dark matter halos being determined by a single parameter κ. Therefore, although lying beyond the range of the present letter, the in-depth comparison of the theoretical predictions of the EiBI gravity model with the galactic and extra galactic scale observations yield tight constraints on the numerical values of κ, thus leading to a direct viability test of the theory. Work along these lines is currently underway.

Acknowledgments. FSNL acknowledges financial support of the Fundação para a Ciência e Tecnologia through the grants CERN/FP/123615/2011 and CERN/FP/123618/2011. SVS acknowledges financial support of the Russian Foundation for Basic Research through grants No. 11-02- 01162 and 13-02-12093 .

[1] M. Aguilar et al. [AMS Collaboration], Phys. Rev. Lett. **110**, 141102 (2013).

[2] H. -B. Jin, Y. -L. Wu and Y. -F. Zhou, arXiv:1304.1997 (2013).

[3] J. Binney and S. Tremaine, *Galactic dynamics*, Princeton University Press, Princeton (1987); M. Persic, P. Salucci and F. Stel, Month. Not. R. Astron. Soc. **281**, 27 (1996); A. Borriello and P. Salucci, Month. Not. R. Astron. Soc. **323**, 285 (2001); P. Salucci et al., Month. Not. Roy. Astron. Soc. **378**, 41 (2007); H. J. de Vega, P. Salucci, and N. G. Sanchez, New Astronomy **17**, 653 (2012).

[4] O. Bertolami, C. G. Boehmer, T. Harko and F. S. N. Lobo, Phys. Rev. D **75**, 104016 (2007); T. Harko and F. S. N. Lobo, Eur. Phys. J. C **70**, 373 (2010).

[5] A. S. Eddington, *The mathematical theory of relativity*, (Cambridge University Press, Cambridge, 1924).

[6] M. Born and L. Infeld, Proc. R. Soc. A **144**, 425 (1934).

[7] M. Bañados and P. G. Ferreira, Phys. Rev. Lett. **105**, 011101 (2010).

[8] T. Delsate and J. Steinhoff, Phys. Rev. Lett. **109**, 021101 (2012).

[9] P. Pani, T. Delsate, and V. Cardoso, Phys. Rev. D **85**, 084020 (2012).

[10] J. H. C. Scargill, M. Bañados, and P. G. Ferreira, Phys. Rev. D **86**, 103533 (2012).

[11] P. Pani and T. P. Sotiriou, Phys. Rev. Lett. **109**, 251102 (2012).

[12] P. Pani, V. Cardoso, and T. Delsate, Phys. Rev. Lett. **107**, 031101 (2011); P. P. Avelino, Phys. Rev. D **85**, 104053 (2012); I. L. J. Casanellas, P. Pani, and V. Cardoso, Astrophys. J. **745**, 15 (2012).

[13] Y.-H. Sham, L.-M. Lin, and P. T. Leung, Phys. Rev. D **86**, 064015 (2012); Y.-H. Sham, P. T. Leung, L.-M. Lin, Phys. Rev. D **87**, 061503(R) (2013).

[14] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, Oxford, Pergamon Press (1975).

[15] T. Matos, F. S. Guzman and D. Nunez, Phys. Rev. D **62**, 061301 (2000); U. Nucamendi, M. Salgado and D. Sudarsky, Phys. Rev. D **63**, 125016 (2001); M. K. Mak and T. Harko, Phys. Rev. D **70**, 024010 (2004); C. G. Boehmer, T. Harko, and F. S. N. Lobo, Astropart. Phys. **29**, 386 (2008); T. Harko, Phys. Rev. D **81**, 084050 (2010).

[16] P. Salucci and M. Persic, in *Dark and visible matter in galaxies*, eds. M. Persic and P. Salucci, ASP Conference Series **117**, 1 (1997).

[17] S. Chandrasekhar, *An introduction to the study of stellar structure*, New York, Dover Publications (1957); G. P. Horedt, *Polytropes: applications in astrophysics and related fields*, Dordrecht, Boston, Kluwer Academic Publishers (2004).

[18] C. G. Boehmer and T. Harko, JCAP **0706**, 025 (2007); T. Harko, Month. Not. Roy. Astron. Soc. **413**, 3095 (2011); T. Harko, JCAP **1105**, 022 (2011).